arXiv:1406.3503v1 [math.AG] 13 Jun 2014

The First and Second Most Symmetric Nonsingular Cubic Surfaces

H. Kaneta, S. Marcugini and F.Pambianco

June 16, 2014

Abstract

The first and second most symmetric nonsingular cubic surfaces are $x^3+y^3+z^3+t^3=0$ and $x^2y+y^2z+z^2t+t^2x=0$, respectively.

1 Introduction

Throughout this paper k stands for an algebraically closed field of characteristic zero. Let C_d be a nonsingular curve of degree $d \geq 4$ in the two-dimensional projective space. As is well known, its automorphism group $\mathrm{Aut}(C_d)$ is a finite subgroup of $PGL_3(k)$ such that $|\mathrm{Aut}(C_d)| \leq 42d(d-3)$. In fact the maximum value β_d of $|\mathrm{Aut}(C_d)|$ is equal to 168, 360 or $6d^2$, and attained by the Klein quartic curve, the Wiman sextic curve or the Fermat curve according as $d=4$, $d=6$ or $d \notin \{4,6\}$[5]. Moreover the Klein quartic and the Wiman sextic give rise to highly symmetric MDS codes [2].

Let S_d be a nonsingular surface of degree $d \geq 3$ in the three-dimensional projective space. Its automorphism group $\mathrm{Aut}(S_d)$ is a finite subgroup of $PGL_4(k)$ if $d \neq 4$, can be an infinite group if $d=4$ [4]. Highly symmetric surfaces may be both of theoretical and of practical interest. Hosoh has completed the classification of the automorphism groups of nonsingular cubic surfaces [1]. According to his classification the maximum and the second maximum of $|\mathrm{Aut}(S_3)|$ is attained by the semidirect product $(\mathbf{Z}_3)^3 \times_s \mathbf{S}_4$ and $\mathbf{S}_5$, respectively. It may happen that $|\mathrm{Paut}(S_d)|$, the order of the projective automorphism group of S_d, has the finite upper bound for every $d \geq 3$. In this paper we will specify S_3 such that $\mathrm{Aut}(S_3)$ is isomorphic to $(\mathbf{Z}_3)^3 \times_s \mathbf{S}_4$ or $\mathbf{S}_5$.

$M_{m,n}(k)$ stands for the set of all $m \times n$ matrices of entries in k. By definition $M_n(k) = M_{n,n}(k)$, $GL_n(k) = \{A = [a_{ij}] \in M_n(k) \ : \ \det A \neq 0\}$, and $PGL_n(k) = GL_n(k)/(E_n)$, where (E_n) is the subgroup $\{\lambda E_n \ : \ \lambda \in k^*\}$ (E_n is the unit matrix in $GL_n(k)$). The coset $A(E_n)$ containing an $A \in GL_n(k)$ will be denoted (A). We denote by $k[x]$ the k-algebra of polynomials in $x=[x_1,...,x_n]$ over k. For an $A \in GL_n(k)$ and $f \in k[x]$ we define a polynomial $f_A \in k[x]$ to be $f_A(x) = f(\sum \alpha_{1j}x_j, ..., \sum \alpha_{nj}x_j)$, where $A^{-1} = [\alpha_{ij}]$. As is well known, the map $T_A : k[x] \to k[x]$ assigning f_A to f is a k-algebra isomorphism of $k[x]$ such that $T_AT_B = T_{AB}$, that is, $(f_B)_A = f_{AB}$. $\mathbf{S}_n$ denotes the symmetric group, the group of all permutations of n elements.

A homogeneous polynomial f of degree $d \geq 1$ defines a projective algebraic set $V(f) = \{(a) \in P^{n-1} \ : \ f(a)=0\}$ of an $(n-1)-$dimensional projective space P^{n-1} over k. $V(f)$ is called a hypersurface of degree d. If $n=4$, $d=3$, and $V(f)$ is nonsingular, the automorphism group $\mathrm{Aut}(V(f))$ of the hypersurface $V(f)$ is a finite subgroup of $PGL_4(k)$ consisting of $(A) \in PGL_4(k)$ such that $f_A \sim f$ [4]. Let $a=[a_1,...,a_n] \in k^n$, $(a) \in V(f)$, and $A \in GL_n(k)$. Then (a) is a singular point of $V(f)$ if $f_{x_i}(a)=0$ for all i. If (a) is a nonsingular point of $V(f)$, $V(\sum_{i=1}^n \gamma_i x_i)$ is the tangent plane to $V(f)$ at (a), where

$\gamma_i = f_{x_i}(a)$. Clearly $(A) : V(f) \to V(f_A)$ is a bijection, and if $b = Aa$ with $(a) \in V(f)$, then $(f_A)_{x_j}(b) = \sum_{i=1}^{n} f_{x_i}(a)\alpha_{ij}$, where $A^{-1} = [\alpha_{ij}]$. Consequently $(b) = (A)(a)$ is a nonsingular point of $V(f_A)$ if and only if (a) is a nonsingular point of $V(f)$, and the tangent plane of $V(f_A)$ at (b) coincides with $(A)V(\sum_{i=1}^{n} \gamma_i x_i)$. In particular if (a) is a nonsingular point of $V(f)$, $f_A \sim f$ and $(A)(a) = (a)$, then $[f_{x_1}(a), ..., f_{x_n}(a)]A \sim [f_{x_1}(a), ..., f_{x_n}(a)]$.

In §1 it will be shown that the Fermat surface $S = V(x^3 + y^3 + z^3 + t^3)$ is the unique cubic nonsingular surface, up to projective equivalence, such that $\mathrm{Aut}(S)$ is isomorphic to $(\mathbf{Z}_3)^3 \times_s \mathbf{S}_4$. In §2 it will be shown that a surface $S' = V(x^2t + y^2z + z^2t + t^2x)$ is the unique cubic nonsingular surface such that $\mathrm{Aut}(S')$ is isomorphic to $\mathbf{S}_5$.

2 ${\mathbf{Z}_3}^3 \times_s \mathbf{S}_4$-invariant nonsingular cubic surfaces

Let $\omega \in k^*$ be of order three. Any subgroup of $PGL_4(k)$ isomorphic to $\mathbf{Z}_3$ is conjugate to $\langle(\mathrm{diag}[\omega, 1, 1, 1])\rangle$ or $\langle(\mathrm{diag}[\omega, \omega^2, 1, 1])\rangle$. It can be verified easily that $X = [x_{ij}] \in GL_4(k)$ satisfies $\mathrm{diag}[\omega, 1, 1, 1]X \sim X\mathrm{diag}[\omega, 1, 1, 1]$ (resp. $\mathrm{diag}[\omega, \omega^2, 1, 1]X \sim X\mathrm{diag}[\omega, \omega^2, 1, 1]$) if and only if $x_{1i} = x_{i1} = 0$ for all $i \in [2, 4]$ (resp. $x_{1i} = x_{i1} = x_{2j} = x_{j2} = 0$ for all $i \in [2, 4]$ and all $j \in [3, 4]$). Consequently a subgroup of $PGL_4(k)$ isomorphic to $(\mathbf{Z}_3)^2$ is conjugate to $\langle(A), (B)\rangle$, where $\mathrm{ord}(A) = \mathrm{ord}(B) = 3$ and $(B^j) \in \langle(A)\rangle$ if and only if $j \in 3\mathbf{Z}$. We may assume that $A = \mathrm{diag}[\omega, \omega^2, 1, 1]$ or $A = \mathrm{diag}[\omega, 1, 1, 1]$ and that $B = \mathrm{diag}[b_1, b_2, b_3, b_4]$. Assume first that $A = \mathrm{diag}[\omega, \omega^2, 1, 1]$. If $|\{b_1, b_2, b_3, b_4\}| = 3$, then we may assume that B is equal to one of $S - \{A\}$, where

$$\begin{aligned} S \;=\; & \{\mathrm{diag}[1, 1, \omega, \omega^2], \mathrm{diag}[1, \omega, 1, \omega^2], \mathrm{diag}[1, \omega, \omega^2, 1], \mathrm{diag}[\omega, 1, 1, \omega^2], \\ & \mathrm{diag}[\omega, 1, \omega^2, 1], \mathrm{diag}[\omega, \omega^2, 1, 1]\}, \end{aligned}$$

Unless $B = \mathrm{diag}[1, 1, \omega, \omega^2]$, there exist integers i and j such that (A^iB^j) is equal to $(\mathrm{diag}[1, 1, \omega, 1])$ or $(\mathrm{diag}[1, 1, 1, \omega])$. So $\langle(A), (B)\rangle$ is conjugate to

$$\langle(\mathrm{diag}[\omega, \omega^2, 1, 1]), (\mathrm{diag}[1, 1, \omega, \omega^2])\rangle \;\; or \;\; \langle(\mathrm{diag}[\omega, \omega^2, 1, 1]), (\mathrm{diag}[1, 1, \omega, 1])\rangle,$$

provided $|\{b_1, b_2, b_3, b_4\}| = 3$. Clearly $\langle(A), (B)\rangle$ is conjugate to

$$\langle(\mathrm{diag}[\omega, \omega^2, 1, 1]), (\mathrm{diag}[1, 1, \omega, 1)\rangle \;\; or \;\; \langle(\mathrm{diag}[\omega, 1, 1, 1]), (\mathrm{diag}[1, 1, \omega, 1])\rangle,$$

provided $|\{b_1, b_2, b_3, b_4\}| = 2$. Assume secondly that $A = \mathrm{diag}[\omega, 1, 1, 1]$. According as $|\{b_1, b_2, b_3, b_4\}|$ is two or three, we may assume B is equal to one of $T - \{A\}$ or S, where

$$T \;=\; \{\mathrm{diag}[\omega, 1, 1, 1], \mathrm{diag}[1, \omega, 1, 1], \mathrm{diag}[1, 1, \omega, 1], \mathrm{diag}[1, 1, 1, \omega]\}.$$

Thus a subgroup of $PGL_4(k)$ isomorphic to $(\mathbf{Z}_3)^2$ is conjugate to one of

$$\langle(\mathrm{diag}[\omega, \omega^2, 1, 1]), (\mathrm{diag}[1, 1, \omega, \omega^2)]\rangle,$$

$$\langle(\mathrm{diag}[\omega, \omega^2, 1, 1]), (\mathrm{diag}[1, 1, \omega, 1])\rangle,$$

$$\langle(\mathrm{diag}[\omega, 1, 1,]), (\mathrm{diag}[1, \omega, 1, 1])\rangle.$$

Even though the following lemma may be known, we shall give a proof for the sake of completeness.

Lemma 2.1. *A subgroup of $PGL_4(k)$ isomorphic to $(\mathbf{Z}_3)^3$ is conjugate to*

$$G_{27} = \langle(\mathrm{diag}[\omega, 1, 1, 1]), (\mathrm{diag}[1, \omega, 1, 1]), (\mathrm{diag}[1, 1, \omega, 1])\rangle.$$

Proof. Let $G = \langle (A), (B), (C) \rangle$ be a subgroup of $PGL_4(k)$ isomorphic to $(\mathbf{Z}_3)^3$. Since $\langle (A), (B) \rangle$ is isomorphic to $(\mathbf{Z}_3)^2$, we may assume one of the following three cases: 1) $A = \mathrm{diag}[\omega, \omega^2, 1, 1]$, $B = \mathrm{diag}[1, 1, \omega, \omega^2]$. 2) $A = \mathrm{diag}[\omega, \omega^2, 1, 1]$, $B = \mathrm{diag}[1, 1, \omega, 1]$. 3) $A = \mathrm{diag}[\omega, 1, 1, 1]$, $B = \mathrm{diag}[1, \omega, 1, 1]$. In any case we may assume $C = \mathrm{diag}[c_1, c_2, c_3, c_4] \neq E_4$ with $c_i^3 = 1$ but $(C) \notin \langle (A), (B) \rangle$. Let $m = |\{c_1, c_2, c_3, c_4\}|$. Assume the case 1). If $m = 2$, then $G = G_{27}$. If $m = 3$, then we may assume C is equal to one of $S - \{A, B\}$ so that G contains one of T, hence $G = G_{27}$. Assume the case 2). If $m = 2$, then we may assume C is one of $T - \{B\}$, hence $G = G_{27}$. If $m = 3$, then we may assume C is one of $S - \{\mathrm{diag}[1, \omega, 1, \omega^2], \mathrm{diag}[\omega, 1, 1, \omega^2], A\}$, hence $G = G_{27}$. Note that $(\mathrm{diag}[1, \omega, 1, \omega^2])$, $(\mathrm{diag}[\omega, 1, 1, \omega^2]) \in \langle (A), (B) \rangle$. Assume the case 3). If $m = 2$, then we may assume C is one of $T - \{A, B\}$, hence $G = G_{27}$. If $m = 3$, then we may assume C is one of $S - \{\mathrm{diag}[\omega, \omega^2, 1, 1]\}$, hence $G = G_{27}$.

The canonical group representation $\hat{}\colon \mathbf{S}_4 \to GL_4(k)$ of $\mathbf{S}_4$ is the one such that $\hat{\sigma}x = y$ with $y_i = x_{\sigma^{-1}(i)}$ for any column vector $x \in k^4$. Clearly this representation is an isomorphism. Let $r \geq 2$ be an integer, $\delta \in k^*$ be of order r,

$$D(r) = \{\mathrm{diag}[\lambda_1, \lambda_2, \lambda_3, \lambda_4] \in GL_4(k) \ : \ \lambda_1^r = \lambda_2^r = \lambda_3^r = \lambda_4^r\}.$$

The factor group $(D(r)) = D(r)/k^*E_4$ is isomorphic to $\{\mathrm{diag}[\delta^i, \delta^j, \delta^\ell, 1] \ : \ i, j, \ell \in [0, r-1]\}$ which is isomorphic to $(\mathbf{Z}_r)^3$. A map $\varphi_\sigma : (D(r)) \to (D(r))$ defined by $\varphi_\sigma((A)) = (\hat{\sigma}A\hat{\sigma}^{-1})$ is a group automorphism such that $\varphi_{\sigma\tau} = \varphi_\sigma \circ \varphi_\tau$. The factor group $D(r)\hat{\mathbf{S}}_4/k^*E_4$ is isomorphic to the semidirect product $(D(r)) \times_s \mathbf{S}_4$ such that $((A), \sigma)((A'), \sigma') = ((A)\varphi_\sigma(A'), \sigma\sigma')$. Clearly $(D(r)) \times_s \mathbf{S}_4$ is isomorphic to $(\mathbf{Z}_r)^3 \times_s \mathbf{S}_4$. We note that $\hat{\sigma}\mathrm{diag}[a_1, a_2, a_3, a_4]\hat{\sigma}^{-1} = \mathrm{diag}[b_1, b_2, b_3, b_4]$, where $b_i = a_{\sigma^{-1}(i)}$. We may skip the proof of the following lemma [6].

Lemma 2.2. *If $r \geq 3$, then the projective automorphism group of the surface $V(x^r + y^r + z^r + t^r)$ is $D(r)\hat{\mathbf{S}}_4/k^*E_4$, which can be identified with $(D(r)) \times_s \mathbf{S}_4$.*

Lemma 2.3. *Any $(\mathbf{Z}_3)^3$-invariant nonsingular cubic surface is projectively equivalent to $V(x^3 + y^3 + z^3 + t^3)$. In particular any $(\mathbf{Z}_3)^3 \times_s \mathbf{S}_4$-invariant nonsingular cubic surface is projectively equivalent to $V(x^3 + y^3 + z^3 + t^3)$, and its automorphism group is conjugate to $(D(3)) \times_s \mathbf{S}_4$.*

Proof. By Lemma 1.1 it suffices to show that any G_{27}-invariant nonsingular cubic surface is $V(ax^3 + by^3 + cz^3 + dt^3)$, where a, b, c and d are nonzero constants. Let $V(f)$ be a G_{27}-invariant nonsingular cubic surface, where the homogeneous polynomial $f(x, y, z, t)$ of degree three has the form

$$\begin{aligned} &a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3 \\ &+x^2(b_{12}y + b_{13}z + b_{14}t) + y^2(b_{21}x + b_{23}z + b_{24}t) + z^2(b_{31}x + b_{32}y + b_{34}t) \\ &+t^2(b_{41}x + b_{42}y + b_{43}z) + c_1yzt + c_2xzt + c_3xyt + c_4xyz. \end{aligned}$$

Let

$$A_1 = \mathrm{diag}[\omega, 1, 1, 1], \ A_2 = \mathrm{diag}[1, \omega, 1, 1], \ A_3 = \mathrm{diag}[1, 1, \omega, 1].$$

It is evident that $\mathrm{ord}(A_i) = 3$. Since G_{27} contains (A_i) $(i \in [1, 4])$, $f_{A_i^{-1}}$ is equal to one of $\{f, \ \omega f, \ \omega^2 f\}$. If $f_{A_1^{-1}}$ is equal to ωf or $\omega^2 f$, then f is divisible by x, hence $V(f)$ is singular. Assume $f_{A_1^{-1}} = f$. Then

$$f(x, y, z, t) = a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3 + y^2(b_{23}z + b_{24}t) + z^2(b_{32}y + b_{34}t) + t^2(b_{42}y + b_{43}z) + c_1yzt.$$

Unless $f_{A_2^{-1}} = f$, $V(f)$ is singular. Therefore $f(x, y, z, t) = a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3 + b_{34}z^2t + b_{43}t^2z$. Unless $f_{A_3^{-1}} = f$, $V(f)$ is singular. Thus $f(x, y, z, t) = a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3$. Now $V(f)$ is nonsingular if and only if $a_1a_2a_3a_4 \neq 0$.

3 $\mathbf{S}_5$-invariant nonsingular cubic surfaces

A subgroup of $PGL_4(k)$ isomorphic to the symmetric group $\mathbf{S}_5$ is one of three groups $C_{5!}\mathrm{I}$, $C_{5!}\mathrm{II}$, $C_{5!}\mathrm{III}$ up to conjugacy [3]. We denote these groups by $G(1)$, $G(2)$ and $G(3)$, respectively. There exist group isomorphisms $\varphi_i : \mathbf{S}_5 \to G(i)$ such that $(E_1) = \varphi_i((123))$, $(E_2) = \varphi_i((12)(34))$, $(E_3) = \varphi_i((12)(45))$ and $(F) = \varphi_i((12))$ generate $G(i)$ ($i \in [1,3]$) (cf [7, (2.14) in chap.3]). Note that $(123) = (12)(23)$. Let $\omega = \frac{-1+i\sqrt{3}}{2}$. Then $G(1) = \langle (E_1), (E_2), (E_3), (F) \rangle$, where

$$E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \omega & 0 \\ 0 & 0 & 0 & \omega^2 \end{bmatrix}, \ E_2 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ 0 & \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \\ 0 & \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \end{bmatrix}, \ E_3 = \begin{bmatrix} \frac{1}{4} & -\frac{\sqrt{15}}{4} & 0 & 0 \\ \frac{\sqrt{15}}{4} & \frac{1}{4} & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix},$$

$$F = \frac{1+i}{\sqrt{2}} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

Lemma 3.1. *The $G(1)$-invariant nonsingular cubic surface is $V(f)$, where*

$$f(x,y,z,t) = 3\sqrt{15}x^3 + 10(y^3+z^3+t^3) - 3\sqrt{15}xy^2 - 6(\sqrt{15}x+5y)zt.$$

Moreover, $\mathrm{Aut}(V(f)) = G(1)$.

Proof. Let $F' = \frac{\sqrt{2}}{1+i}F$. Assume that $V(f)$ is a $G(1)$-invariant nonsingular cubic surface, where $f(x,y,z,t)$ is a cubic homogeneous polynomial of the form as in the proof of Lemma 1.3. Since $E_1^3 = E_4$, $f_{E_1^{-1}} = \omega^i f$ for some $i \in [0,2]$. If $i = 1$, then $f(x,y,z,t) = b_{13}x^2z + b_{23}y^2z + b_{34}z^2t + t^2(b_{41}x + b_{42}y) + c_4xyz$. Since $f_{F'^{-1}} \sim f$, we see $b_{13} = b_{23} = 0$, hence $V(f)$ is singular at $(1,0,0,0)$. Similarly $i \neq 2$, namely $i = 0$, so that $f(x,y,z,t) = a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3 + b_1x^2y + b_2y^2x + c_1yzt + c_2xzt$. We have $a_3a_4 \neq 0$, for $V(f)$ is nonsingular. If $a_1 = a_2 = 0$, the condition $f_{E_3^{-1}} \sim f$ yields $b_1 = b_2 = 0$, hence $V(f)$ is singular. Thus $[a_1, a_2] \neq [0,0]$. Therefore $f_{F'^{-1}} = f$, hence $a_3 = a_4$. Now $f_{E_3^{-1}} = f$, and $f_{E_3^{-1}} = -f$ does not hold. The condition $f_{E_3^{-1}} = f$ is equivalent to $c_2 = \sqrt{15}c_1/5$ and

$$\begin{bmatrix} -65 & 15\sqrt{15} & \sqrt{15} & -15 \\ 15\sqrt{15} & -63 & 15 & \sqrt{15} \\ 3\sqrt{15} & 45 & -93 & 13\sqrt{15} \\ -45 & 3\sqrt{15} & 13\sqrt{15} & -35 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ b_1 \\ b_2 \end{bmatrix} = 0.$$

The general $[a_1, a_2, b_1, b_2]$ satisfying the second condition above is $\lambda[3\sqrt{15}, 10, 0, -3\sqrt{15}]$, for the rank of the above matrix is equal to three. Consequently we may assume $f(x,y,z,t) = 3\sqrt{15}x^3 + 10y^3 + a_3(z^3+t^3) - 3\sqrt{15}xy^2 + c_1(yzt + \frac{\sqrt{15}}{5}xzt)$. Now the condition $f_{E_3^{-1}} \sim f$ implies $f_{E_3^{-1}} = f$. Using the formula $(a+b+c)^3 = a^3+b^3+c^3+3a^2(b+c)+3b^2(a+c)+$

$3c^2(a+b)+6abc$, we can write $f_{E_3^{-1}}(x,y,z,t)$ as

$$
\begin{aligned}
&3\sqrt{15}x^3+(-\frac{10}{27}+\frac{16}{27}a_3-\frac{4}{27}c_1)y^3+(\frac{80}{27}+\frac{7}{27}a_3-\frac{4}{27}c_1)z^3+(\frac{80}{27}+\frac{7}{27}a_3-\frac{4}{27}c_1)t^3\\
&+y^2\{(-\frac{\sqrt{15}}{3}+\frac{4\sqrt{15}}{45}c_1)x+(\frac{60}{27}+\frac{12}{27}a_3+\frac{6}{27}c_1)(z+t)\}\\
&+z^2\{(-\frac{4\sqrt{15}}{3}-\frac{2\sqrt{15}}{45}c_1)x+(-\frac{120}{27}+\frac{30}{27}a_3+\frac{6}{27}c_1)y+(\frac{240}{27}-\frac{6}{27}a_3+\frac{6}{27}c_1)t\}\\
&+t^2\{(-\frac{4\sqrt{15}}{3}-\frac{2\sqrt{15}}{45}c_1)x+(-\frac{120}{27}+\frac{30}{27}a_3+\frac{6}{27}c_1)y+(\frac{240}{27}-\frac{6}{27}a_3+\frac{6}{27}c_1)z\}\\
&+(-\frac{240}{27}-\frac{48}{27}a_3+\frac{3}{27}c_1)yzt+(-\frac{8\sqrt{15}}{3}+\frac{\sqrt{15}}{9}c_1)xzt\\
&+(\frac{4\sqrt{15}}{3}+\frac{2\sqrt{15}}{45}c_1)xyt+(\frac{4\sqrt{15}}{3}+\frac{2\sqrt{15}}{45}c_1)xyz.
\end{aligned}
$$

Now $a_3=10$, $c=-30$ and $f_{E_3^{-1}}=f$ holds, as desired. We can show that $V(f)$ is nonsingular.

Let $\varepsilon=-\frac{\sqrt{5}-1}{4}+i\frac{\sqrt{10+2\sqrt{5}}}{4}$, hence $\mathrm{ord}(\varepsilon)=5$, $\alpha=\frac{-\sqrt{5}+1}{2}$, $\beta=\alpha^2$, $\gamma=-\alpha$, and let

$$
H=\begin{bmatrix}\varepsilon^4&0&0&0\\0&\varepsilon^2&0&0\\0&0&\varepsilon&0\\0&0&0&\varepsilon^3\end{bmatrix},\ I=\begin{bmatrix}1&\alpha&\beta&\gamma\\\alpha&\beta&\gamma&1\\\beta&\gamma&1&\alpha\\\gamma&1&\alpha&\beta\end{bmatrix}.
$$

Lemma 3.2. *(H) and (I) generate in $PGL_4(k)$ a group $G(1)'$ conjugate to $G(1)$. The $G(1)'$-invariant nonsingular cubic surface is $V(x^2y+y^2z+z^2t+t^2x)$. Moreover* $\mathrm{Aut}(V(x^2y+y^2z+z^2t+t^2x))=G(1)'$.

Proof. The transpositions $(i\ \ i+1)$ $(i\in[1,4])$ generate $\mathbf{S}_5$, while $(12345)(j\ \ j+1)(12345)^{-1}=(j+1\ \ j+2)$ $(j\in[1,3])$. Therefore (12345) and (12) generate $\mathbf{S}_5$. Clearly $(123)(34)(45)=(12345)$. In view of the group isomorphism $\varphi_1: S_5\to G(1)$, $\varphi_1((12345))=(E_1FE_2FE_3)$ and $\varphi_1((12))=(F)$. Let

$$
K=E_1FE_2FE_3=\begin{bmatrix}-\frac{1}{4}&\frac{\sqrt{15}}{4}&0&0\\-\frac{\sqrt{15}}{12}&-\frac{1}{12}&\frac{2}{3}&\frac{2}{3}\\\frac{\sqrt{15}\omega}{6}&\frac{\omega}{6}&\frac{2\omega}{3}&-\frac{\omega}{3}\\\frac{\sqrt{15}\omega^2}{6}&\frac{\omega^2}{6}&-\frac{\omega^2}{3}&\frac{2\omega^2}{3}\end{bmatrix},\ T=\begin{bmatrix}1&0&0&0\\0&1&0&0\\0&0&1&-1\\0&0&1&1\end{bmatrix}.
$$

Now

$$
T^{-1}FT=\begin{bmatrix}1&0&0&0\\0&1&0&0\\0&0&1&0\\0&0&0&-1\end{bmatrix},\ T^{-1}KT=\begin{bmatrix}-\frac{1}{4}&\frac{\sqrt{15}}{4}&0&0\\-\frac{\sqrt{15}}{12}&-\frac{1}{12}&\frac{4}{3}&0\\-\frac{\sqrt{15}}{12}&-\frac{1}{12}&-\frac{1}{6}&-i\frac{\sqrt{3}}{2}\\-i\frac{\sqrt{5}}{4}&-i\frac{\sqrt{3}}{12}&-i\frac{\sqrt{3}}{6}&-\frac{1}{2}\end{bmatrix},
$$

$\det(T^{-1}KT-\lambda E_4)=\lambda^4+\lambda^3+\lambda^2+\lambda+1=\Pi_{i=1}^4(\lambda-\varepsilon^i)$.

If λ is an eigenvalue of $T^{-1}KT$ and $x\in k^4$ satisfies $(T^{-1}KT-\lambda E_4)x=0$, then

$$
[x_1,x_2,x_3,x_4]=x_1[1,\ (1+4\lambda)\frac{\sqrt{15}}{15},\ (1+\lambda+3\lambda^2)\frac{\sqrt{15}}{15},\ i(-\lambda^4+\lambda^3)\frac{\sqrt{3}}{5}].
$$

So $S^{-1}T^{-1}KTS = \mathrm{diag}[\varepsilon^4, \varepsilon^2, \varepsilon, \varepsilon^3]$, where

$$S = \begin{bmatrix} 1 & 1 & 1 & 1 \\ (1+4\varepsilon^4)\frac{\sqrt{15}}{15} & (1+4\varepsilon^2)\frac{\sqrt{15}}{15} & (1+4\varepsilon)\frac{\sqrt{15}}{15} & (1+4\varepsilon^3)\frac{\sqrt{15}}{15} \\ (1+\varepsilon^4+3\varepsilon^3)\frac{\sqrt{15}}{15} & (1+\varepsilon^2+3\varepsilon^4)\frac{\sqrt{15}}{15} & (1+\varepsilon+3\varepsilon^2)\frac{\sqrt{15}}{15} & (1+\varepsilon^3+3\varepsilon)\frac{\sqrt{15}}{15} \\ i(-\varepsilon+\varepsilon^2)\frac{\sqrt{3}}{5} & i(-\varepsilon^3+\varepsilon)\frac{\sqrt{3}}{5} & i(-\varepsilon^4+\varepsilon^3)\frac{\sqrt{3}}{5} & i(-\varepsilon^2+\varepsilon^4)\frac{\sqrt{3}}{5} \end{bmatrix}$$

so that $\det S = -i\frac{8\sqrt{3}}{25}\varepsilon^3(\varepsilon-1)^3(3\varepsilon^3+6\varepsilon^2+4\varepsilon+2)$. We denote the (i,j)-cofactor of $S = [s_{ij}]$ by $\tilde{s}_{ij}$. Note that $S^{-1}\mathrm{diag}[1,1,1,-1]S = E_4 + S^{-1}\mathrm{diag}[0,0,0,-2]S$ and that the i-th row of $S^{-1}\mathrm{diag}[0,0,0,-2]S$ is equal to $-2\tilde{s}_{4i}/\det S \times$(4-th row of S). Denote $TS\mathrm{diag}[1,\varepsilon^2,\varepsilon^3,\varepsilon]$ by S'. By computation we get the components of $(3\varepsilon^3+6\varepsilon^2+4\varepsilon+2)S'^{-1}FTS$' as follows.

$$\begin{bmatrix} 2\varepsilon^3+4\varepsilon^2+3\varepsilon+1 & 2\varepsilon^4+\varepsilon^3+2 & -\varepsilon^4+\varepsilon^2 & \varepsilon^3+2\varepsilon^2+2\varepsilon \\ 2\varepsilon^4+\varepsilon^3+2 & -\varepsilon^4+\varepsilon^2 & \varepsilon^3+2\varepsilon^2+2\varepsilon & 2\varepsilon^3+4\varepsilon^2+3\varepsilon+1 \\ -\varepsilon^4+\varepsilon^2 & \varepsilon^3+2\varepsilon^2+2\varepsilon & 2\varepsilon^3+4\varepsilon^2+3\varepsilon+1 & 2\varepsilon^4+\varepsilon^3+2 \\ \varepsilon^3+2\varepsilon^2+2\varepsilon & 2\varepsilon^3+4\varepsilon^2+3\varepsilon+1 & 2\varepsilon^4+\varepsilon^3+2 & -\varepsilon^4+\varepsilon^2 \end{bmatrix}.$$

Moreover, since $(2\varepsilon^3+4\varepsilon^2+3\varepsilon+1)^{-1} = (7\varepsilon^3+4\varepsilon^2+\varepsilon+8)/5$, we can verify

$$\begin{aligned} &(2\varepsilon^3+4\varepsilon^2+3\varepsilon+1)^{-1}(2\varepsilon^4+\varepsilon^3+2) = \varepsilon^3+\varepsilon^2+1 = \alpha, \\ &(2\varepsilon^3+4\varepsilon^2+3\varepsilon+1)^{-1}(-\varepsilon^4+\varepsilon^2) = \varepsilon^3+\varepsilon^2+2 = \alpha^2, \\ &(2\varepsilon^3+4\varepsilon^2+3\varepsilon+1)^{-1}(\varepsilon^3+2\varepsilon^2+2\varepsilon) = -\varepsilon^3-\varepsilon^2-1 = -\alpha. \end{aligned}$$

We have shown that $S'^{-1}KS' = H$ and $(S'^{-1}FS') = (I)$.

Finally we shall show that a cubic homogeneous polynomial $f(x,y,z,t)$ of the form as in the proof of Lemma 1.3 such that $f_{H^{-1}} \sim f$, $f_{I^{-1}} \sim f$ and that $V(f)$ is nonsingular is proportional to $x^2y+y^2z+z^2t+t^2x$. Since $H^5 = E_4$, $f_H^{-1} = \varepsilon^i f$ for some $i \in [0,4]$. If $i = 1$, then $f(x,y,z,t) = a_2y^3 + b_{14}x^2t + b_31z^3x + c_1yzt$, hence $V(f)$ is singular at $(0,0,0,1)$. Similarly, unless $i = 0$, $V(f)$ has a singular point. If $i = 0$, then $f(x,y,z,t) = b_1x^2y + b_2y^2z + b_3z^2t + b_4t^2x$. Denoting the coefficients of x^3, y^3, z^3 and t^3 in $f_{I^{-1}}$ by

b'_1, b'_2, b'_3 and b'_4, respectively, and noting $\alpha^2 = \alpha + 1$, we obtain

$$b' = \alpha \begin{bmatrix} 1 & \alpha^3 & -\alpha^4 & \alpha \\ \alpha^3 & -\alpha^4 & \alpha & 1 \\ -\alpha^4 & \alpha & 1 & \alpha^3 \\ \alpha & 1 & \alpha^3 & -\alpha^4 \end{bmatrix} b.$$

Since $b' = 0$ and the rank of the matrix involved is equal to three, it follows that $b_1 = b_2 = b_3 = b_4 \neq 0$. Now let $f = x^2y+y^2z+z^2t+t^2x$. In order to see $f_{I^{-1}} = 5(4\alpha+3)f$, we note that denoting $(x+\alpha y+\beta z+\gamma t)^2(x+\alpha x+\beta y+\gamma z+t)$ by $g(x,y,z,t)$, we have $f_{I^{-1}}(x,y,z,t) = g(x,y,z,t) + g(t,x,y,z) + g(z,t,x,y) + g(y,z,t,x)$.

The group $G(2)$ is generated by (E_1), (E_2), (E_3) and (F), where

$$E_1 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \omega & 0 \\ 0 & 0 & 0 & \omega^2 \end{bmatrix}, \quad E_2 = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 0 & 0 & \sqrt{2} \\ 0 & -1 & \sqrt{2} & 0 \\ 0 & \sqrt{2} & 1 & 0 \\ \sqrt{2} & 0 & 0 & -1 \end{bmatrix}, \quad E_3 = \begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 & 0 \\ \frac{1}{2} & -\frac{\sqrt{3}}{2} & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix},$$

$$F = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{bmatrix}.$$

Lemma 3.3. *There exist no $G(2)$-invariant nonsingular cubic surfaces.*

Proof. Let $f(x,y,z,t)$ be a $G(2)$-invariant cubic homogeneous polynomial of the form as in the proof of Lemma 1.3. Since $F^4 = E_4$, $f_{F^{-1}} = i^j f$ for some $j \in [0,3]$. If $j = 0$ or $j = 2$, then $f = 0$. If $j = 1$, then $f(x,y,z,t)$ has the form

$$\begin{aligned}&a_1(x^3 - iy^3) + a_3(z^3 - it^4)\\ &+x^2(b_{12}y + b_{13}z + b_{14}t) + y^2(ib_{12}x + ib_{14}z - ib_{13}t)\\ &+z^2(b_{31}x + b_{32}y + b_{34}t) + t^2(ib_{32}x - ib_{31}y + ib_{34}z)\\ &+c_1yzt - ic_1xzt + c_3xyt - ic_3xyz.\end{aligned}$$

Since $E_3^2 = E_4$, $f_{E_3^{-1}} = \pm f$. The coefficients of t^3, t^2z, t^2x and t^2y of $f_{E_3^{-1}}$ (resp. $\pm f$) are

$$a_3,\ b_{34},\ \frac{\sqrt{3}}{2}b_{31} + \frac{1}{2}b_{32},\ \frac{1}{2}b_{31} - \frac{\sqrt{3}}{2}b_{32} \text{ (resp. } \mp ia_3,\ \pm ib_{34},\ \pm ib_{32},\ \mp ib_{31}),$$

hence $a_3 = b_{34} = b_{31} = b_{32} = 0$. Consequently $V(f)$ is singular at $(0,0,0,1)$.

Let $\nu_1 = \sqrt{\frac{3}{8}} + i\sqrt{\frac{5}{8}}$ and $\nu_2 = \sqrt{\frac{3}{8}} - i\sqrt{\frac{5}{8}}$ The group $G(3)$ is generated by (E_1), (E_2), (E_3) and (F), where

$$E_1 = \begin{bmatrix} \omega & 0 & 0 & 0\\ 0 & \omega & 0 & 0\\ 0 & 0 & \omega^2 & 0\\ 0 & 0 & 0 & \omega^2 \end{bmatrix},\ E_2 = \frac{1}{\sqrt{3}}\begin{bmatrix} 1 & 0 & 0 & \sqrt{2}\\ 0 & 1 & \sqrt{2} & 0\\ 0 & \sqrt{2} & -1 & 0\\ \sqrt{2} & 0 & 0 & -1 \end{bmatrix},\ E_3 = \begin{bmatrix} 0 & 0 & 0 & \nu_1\\ 0 & 0 & \nu_2 & 0\\ 0 & \nu_1 & 0 & 0\\ \nu_2 & 0 & 0 & 0 \end{bmatrix},$$

$$F = \begin{bmatrix} 0 & 0 & 0 & 1\\ 0 & 0 & -1 & 0\\ 0 & 1 & 0 & 1\\ -1 & 0 & 0 & 0 \end{bmatrix}.$$

Lemma 3.4. *There exist no $G(3)$-invariant nonsingular cubic surfaces. More precisely, there exist no $G(3)$-invariant cubic forms except zero.*

Proof. Let $E_1' = \omega^{-1}E_1$ and $f(x,y,z,t)$ a $G(3)$-invariant cubic form. If $f_{E_1'^{-1}} = \omega f$, then f has the form

$$x^2(b_{13}z + b_{14}t) + y^2(b_{23}z + b_{24}t) + c_3xyt + c_4xyz,$$

hence the condition $f_{F^{-1}} \sim f$ implies $f = 0$. Similarly, if $f_{E_1'^{-1}} = \omega^2 f$, then $f = 0$. Now assume $f_{E_1'^{-1}} = f$, so that $f(x,y,z,t) = a_1x^3 + a_2y^3 + a_3z^3 + a_4t^3 + b_1x^2y + b_2y^2x + b_3z^2t + b_4t^2z$. Since $f_{F^{-1}} = i^j f$ for some $j \in [0,3]$, we obtain

$$\begin{bmatrix} i^j & 1\\ -1 & i^j \end{bmatrix}\begin{bmatrix} a_1\\ a_4 \end{bmatrix} = 0,\ \begin{bmatrix} i^j & -1\\ 1 & i^j \end{bmatrix}\begin{bmatrix} a_2\\ a_3 \end{bmatrix} = 0,\ \begin{bmatrix} i^j & -1\\ 1 & i^j \end{bmatrix}\begin{bmatrix} b_1\\ b_4 \end{bmatrix} = 0,\ \begin{bmatrix} i^j & 1\\ -1 & i^j \end{bmatrix}\begin{bmatrix} b_2\\ b_3 \end{bmatrix} = 0.$$

Therefore, unless $j = 1$ or $j = 3$, $f(x,y,z,t) = 0$. Even if $j = 1$ or $j = 3$, the condition $f_{E_3^{-1}} \sim f$ implies $f = 0$, because $\{i, -i\} \cap \{\nu_1, \nu_2, \nu_1^3, \nu_2^3\} = \emptyset$.

Remark 3.5. *Let $G = \mathrm{Aut}(V(x^2y + y^2z + z^2t + t^2x))$, Σ the field automorphism of $\mathbf{Q}(\varepsilon)$ such that $\varepsilon^\Sigma = \varepsilon^2$.*
(1) *Denoting by e_i be the i-th column of the unit matrix E_4, let $J = [e_2, e_3, e_4, e_1]$. Then $(J) \in G = G(1)'$.*

(2) *Define* $I' \in GL_4(k)$ *as* I*, using* $\alpha' = \frac{\sqrt{5}+1}{2}$*,* $\beta' = \alpha'^2$ *and* $\gamma' = -\alpha'$ *in place of* α*,* β *and* γ*, respectively. Then* $H^\Sigma = H^2$ *and* $I^\Sigma = I'$*, for* $\sqrt{5}^\Sigma = -\sqrt{5}$*. On the other hand* $JHJ^{-1} = H^2$ *and* $JIJ^{-1} = \beta I'$*. Consequently there exists uniquely a group isomorphism* $\psi : \mathbf{S}_5 \to PGL_4(k)$ *such that* $\psi((12345)) = (H^2)$ *and* $\psi((12)) = (I')$*. The projective representations of* $\mathbf{S}_5$ ψ *and* φ_1 *are equivalent.*

References

[1] Hosoh T.: Automorphism groups of cubic surfaces, J. Algebra **192**, 651–677 (1997).

[2] Kaneta, H., Marcugini, S. and Pambianco, F.: On arcs and curves with many automorphisms, Mediterr. J. Math. **2**, 71–102 (2005).

[3] Maschke H.: Bestimmung aller ternären und quaternären Collineationsgruppe, welche mit symmetrischen und alternierenden Buchstabenvertauschungsgruppen holoedrisch isomorph sind, Math. Ann. **51**, 253–298 (1899).

[4] Matsumura H. and Monsky P.: On the automorphisms of hypersurfaces, J. Maht. Kyoto Univ. **3**-3, 347–361 (1964).

[5] Pambianco F.: Characterization of the Fermat curve as the most symmetric nonsingular plane curve, Mathematische Zeitschrift, 2014, DOI 10.1007/s00209-014-1288-4.

[6] Shioda, T.: Arithmetic and geometry of the Fermat curve, Algebraic Seminar (Singapore 1987), World Sci. Publications, Singapore(1988).

[7] Suzuki M.: Group Theory I, Springer 1982.